arXiv:0908.0135v1 [math.OA] 2 Aug 2009

The Rohlin property for automorphisms of the Jiang-Su algebra.

Yasuhiko Sato

Department of Mathematics, Hokkaido University,
Sapporo 060-0810, Japan
e-mail : s053018@math.sci.hokudai.ac.jp

Abstract

For projectionless C^*-algebras absorbing the Jiang-Su algebra tensorially, we study a kind of the Rohlin property for autmorphisms. We show that the crossed products obtained by automorphisms with this Rohlin property also absorb the Jiang-Su algebra tensorially under a mild technical condition on the C^*-algebras.

In particular, for the Jiang-Su algebra we show the uniqueness up to outer conjugacy of the automorphism with this Rohlin property.

Keywords: C^*-algebra, automorphism, Jiang-Su algebra, Rohlin property.

Mathematics Subject Classification 2000: Primary 46L40; Secondary 46L35, 46L80.

1 Introduction

In the classification program established by Elliott, the Jiang-Su algebra $\mathcal{Z}$ is one of the most important C^*-algebras, see [8], [18], and which have been investigated by many people, [4], [6], [10], [29], [33]. Toms and Winter proved that for all approximately divisible C^*-algebras absorb the Jiang-Su algebra tensorially, i.e., $A \cong A \otimes \mathcal{Z}$, [33]. Rørdam showed that the Cuntz semigroup of a $\mathcal{Z}$-absorbing C^*-algebra is almost unperforated [29]. Recently, Winter has shown some criteria for the absorption of the Jiang-Su algebra [35]. For abstract characterization of the Jiang-Su algebra in a streamlined way, we refer to the recent papers by Dadarlat, Rørdam, Toms, and Winter, [6], [30], [35]. H. Lin has shown the classification theorem for a large class of C^*-algebras consisting of generalized dimension drop algebras when they absorb the Jiang-Su algebra tensorially.

In the case of von Neumann algebras Connes defined the Rohlin property for automorphisms, using a partition of unities consisting of projections, and classified automorphisms of injective type II_1 factor up to outer conjugacy [2]. Kishimoto gave a method to prove the Rohlin property for automorphisms of

AF-algebras for classifying automorphisms up to outer conjugacy, based on the Elliott's classification program, [9], [20], [21]. For Kirchberg algebras, Nakamura completely classified autmorphisms with the Rohlin property by their KK-classes up to outer conjugacy [24]. Recently, Matui has classified automorphisms of AH-algebras with real rank zero and slow dimension growth up to outer conjugacy [23]. For finite actions, Izumi defined the Rohlin property and has shown the classification theory [14], [15]. Recently, Izumi, Katsura, and Matui showed classification results for $\mathbb{Z}^2$-actions with the Rohlin property [19], [22], [16].

The aim of this paper is to introduce a kind of the Rohlin property for automorphisms of projectionless C^*-algebras and to give the following two main theorems as follows.

Definition 1.1. Let A be a unital C^*-algebra which has a unique tracial state τ and absorbs the Jiang-Su algebra $\mathcal{Z}$ tensorially, and α be an automorphism of A. We say that α has *the weak Rohlin property*, if for any $k \in \mathbb{N}$ there exist positive elements $f_n \in A_+^1$ such that $(f_n)_n \in A_\infty$, $n \in \mathbb{N}$

$$\begin{aligned}&(\alpha^j(f_n)) \cdot (f_n) = 0, \quad j = 1, 2, ..., k-1,\\ &\tau(1 - \sum_{j=0}^{k-1} \alpha^j(f_n)) \to 0.\end{aligned}$$

Here, we denote by A^∞ the quotient $\ell^\infty(\mathbb{N}, A)/c_0(A)$, and A_∞ the central sequence algebra $A^\infty \cap A'$.

We extend a technical condition called property (SI) to projectionless C^*-algebras in Definition 3.3. Roughly speaking, property (SI) means that if two central sequence of positive elements are given such that one of them is infinitesimally small compared to the other in the sequence algebra, then in fact so in the central sequence algebra.

Theorem 1.2. *Let A be a unital separable projectionless C^*-algebra which has a unique tracial state and absorbs the Jiang-Su algebra tensorially. Suppose that A has property (SI) and α is an automorphism of A with the weak Rohlin property. Then $A \times_\alpha \mathbb{Z}$ also absorbs the Jiang-Su algebra tensorially.*

Theorem 1.3. *Suppose that α and β are automorphisms of the Jiang-Su algebra with the weak Rohlin property. Then α and β are outer conjugate, i.e., there exists an automorphism δ of $\mathcal{Z}$ and a unitary u in $\mathcal{Z}$ such that*

$$\alpha = \mathrm{Ad}\, u \circ \delta \circ \beta \circ \delta^{-1}.$$

For a separable, nuclear C^*-algebra A absorbing the Jiang-Su algebra, Rørdam proved that A is purely infinite if and only if A is traceless in [29], and Nakamura proved that the aperiodicity for automorphisms of purely infinite C^*-algebras coincides with the Rohlin property in [24]. If A is a projectionless C^*-algebra with a unique tracial state constructed in [32], the weak Rohlin property is equivalent to the aperiodicity of the automorphism in the GNS-representation associated with the tracial state. Similar definitions for finite actions, which is called *projection free tracial Rohlin property*, has been defined in [1], [25]. The first main theorem is an adaptation of the result showed by Hirshberg and Winter in [12] to projectionless C^*-algebras. The second main theorem is an adaptation of the result for UHF algebras showed by Kishimoto in [20] to the Jiang-Su algebra.

The paper is organized as follows: In Section 2, we recall the generators of the prime dimension drop algebras discovered by Rørdam and Winter in [30]. In Section 3, for projectionless cases we extend the technical property, which was called property (SI) in [31], to projectionless C^*-algebras. By this property, we can obtain the generators defined in Section 2 . In Section 4, we prove Theorem 1.2. In Section 5, using the weak Rohlin property we show the stability for the automorphisms of the Jiang-Su algebra, and show Theorem 1.3.

Concluding this section, we prepare some notations. When A is a C^*-algebra, we denote by A_{sa} the set of self-adjoint elements of A, A^1 the unit ball of A, A_+ the positive cone of A, $U(A)$ the unitary group of A, $P(A)$ the set of projections of A, $T(A)$ the tracial state space of A. We define an inner automorphism of A by $\operatorname{Ad} u(a) = uau^*$ for $u \in U(A)$ and $a \in A$. We denote by M_n the C^*-algebra of $n \times n$ matrices with complex entries and $e_{i,j}^{(n)}$ the canonical matrix units of M_n, and we set $e_i^{(n)} = e_{i,i}^{(n)}$. We denote by (m, n) the greatest common divisor of m and $n \in \mathbb{N}$.

2 The generators of prime dimension drop algebras

The following argument was given by Rørdam and Winter in [29] and [30]. We would like to begin with some definitions about the generators of prime dimension drop algebras and show Proposition 2.1. We denote by $I(k, k+1)$, $k \in \mathbb{N}$ the prime dimension drop algebra

$$\{f \in C([0,1]) \otimes M_k \otimes M_{k+1};\ f(0) \in M_k \otimes 1_{k+1},\ f(1) \in 1_k \otimes M_{k+1}\},$$

and set the self adjoint unitary

$$u_1 = \sum_{i,j} e_{i,j}^{(k)} \otimes e_{j,i}^{(k)} \in U(M_k \otimes M_k).$$

Define non-unital $*$-homomorphisms $\rho_0 : M_k \otimes M_k \hookrightarrow M_k \otimes M_{k+1}$ by $\rho_0(e_{i,j}^{(k)} \otimes e_{l,m}^{(k)}) = e_{i,j}^{(k)} \otimes e_{l,m}^{(k+1)}$, and $\rho : C([0,1]) \otimes M_k \otimes M_k \hookrightarrow C([0,1]) \otimes M_k \otimes M_{k+1}$ by $\rho(f)(t) = \rho_0(f(t))$, $t \in [0,1]$. Let $u \in U(C([0,1]) \otimes M_k \otimes M_k)$ be such that

$u(0) = 1$ and $u(1) = u_1$, and set

$$\begin{aligned} v &= \sum_{j=1}^{k} e_{1,j}^{(k)} \otimes e_{j,k+1}^{(k+1)}, \\ w(t) &= \rho(u)(t) \oplus \cos^{1/2}(\pi t/2) 1_k \otimes e_{k+1}^{(k+1)}, \\ c_j(t) &= w(t)(e_{1,j}^{(k)} \otimes 1_{k+1}) w^*(t), \quad j = 1, 2, ..., k, \\ s(t) &= \sin(\pi t/2) w(t) v, \quad t \in [0, 1]. \end{aligned}$$

Since $c_j(0) = e_{1,j}^{(k)} \otimes 1_{k+1}$, $c_j(1) = 1_k \otimes e_{1,j}^{(k+1)}$, $s(0) = 0$, and $s(1) = 1_k \otimes e_{1,k+1}^{(k+1)}$, it follows that c_j, $s \in I(k, k+1)$. And we have that

$$\begin{aligned} w^* w(t) &= ww^*(t) = 1_k \otimes (1_{k+1} - e_{k+1}^{(k+1)}) \oplus \cos(\pi t/2) 1_k \otimes e_{k+1}^{(k+1)}, \\ c_i c_j^* &= ww^*w(e_{1,i}^{(k)} e_{j,1}^{(k)} \otimes 1_{k+1}) w^* = \delta_{i,j} c_1^2, \\ \sum_{j=1}^{k} c_j^* c_j &= (w^* w)^2, \\ s^* s(t) &= \sin^2(\pi t/2) 1_k \otimes e_{k+1}^{(k+1)}, \\ c_1 s(t) &= \sin(\pi t/2) w(e_1^{(k)} \otimes 1_{k+1}) w^* w(t) (\sum_{i=1}^{k} e_{1,i}^{(k)} \otimes e_{i,k+1}^{(k+1)}) = s(t). \end{aligned}$$

From these computations, it follows that $\{c_j\}_{j=1}^k \cup \{s\}$ satisfies

$$c_1 \geqq 0, \quad c_i c_j^* = \delta_{i,j} c_1^2,$$

$$\sum_{j=1}^{k} c_j^* c_j + s^* s = 1, \quad c_1 s = s.$$

To be convenient, we denote by $\mathcal{R}_k$ the above relations on generators of a unital C^*-algebra.

Fix a separable infinite-dimensional Hilbert space $\mathcal{H}$, and set

$$\Lambda = \{\{c'_j\}_{j=1}^k \cup \{s'\} \subset B(\mathcal{H})^1; \text{ satisfies } \mathcal{R}_k\} \subset 2^{B(\mathcal{H})^1}.$$

For $\lambda \in \Lambda$, let $c_{j,\lambda} \in \lambda$, $j = 1, 2, ..., k$ and $s_\lambda \in \lambda$ be generators corresponding to c_j, $j = 1, 2, ..., k$, and s on the relations $\mathcal{R}_k$, and define $\tilde{c}_j = \bigoplus_{\lambda \in \Lambda} c_{j,\lambda}$, $\tilde{s} = \bigoplus_{\lambda \in \Lambda} s_\lambda \in B(\bigoplus_{\lambda \in \Lambda} \mathcal{H})$. The set $\{\tilde{c}_j\}_{j=1}^k \cup \{\tilde{s}\}$ satisfies the relations $\mathcal{R}_k$. Let $C^*(\{\tilde{c}_j\}_{j=1}^k \cup \{\tilde{s}\})$ be the C^*-subalgebra of $B(\bigoplus_{\lambda \in \Lambda} \mathcal{H})$ generated by $\{\tilde{c}_j\} \cup \{\tilde{s}\}$. Then, we can identify $C^*(\{\tilde{c}_j\}_{j=1}^k \cup \{\tilde{s}\})$ with the universal C^*-algebra on a set of generators satisfying the relations $\mathcal{R}_k$.

Proposition 2.1 (Proposition 5.1 in [30])**.** *The universal C^*-algebra $C^*(\{\tilde{c}_j\}_{j=1}^k \cup \{\tilde{s}\})$ is isomorphic to $I(k, k+1)$ with $\tilde{c}_j \mapsto c_j$ and $\tilde{s} \mapsto s$.*

Proof. First we show that $C^*(\{c_j\}_j \cup \{s\}) = I(k, k+1)$. Since

$$\sum_{j=1}^{k} c_j^* s s^* c_j + s^* s(t) \quad = \quad \sin^2(\pi t/2)(1_k \otimes 1_{k+1}),$$

and $1_k \otimes 1_{k+1} \in C^*(\{c_j\}_j \cup \{s\})$, we have that $C([0,1]) \otimes 1_k \otimes 1_{k+1} \subset C^*(\{c_j\}_j \cup \{s\})$. By a partition of unity argument on $[0,1]$, it suffices to show that $C^*(\{c_j\}_j \cup \{s\})(i) \cong M_{k+i}$, $i = 0, 1$, and $C^*(\{c_j\}_j \cup \{s\})(t) \cong M_k \otimes M_{k+1}$, $t \in (0,1)$. Since $c_j(0) = e_{1,j}^{(k)} \otimes 1_{k+1}$, $c_j(1) = 1_k \otimes e_{1,j}^{(k+1)}$, $j = 1, 2, \ldots, k$, and $s(1) = 1_k \otimes e_{1,k+1}^{(k+1)}$, it follows that $C^*(\{c_j\}_j \cup \{s\})\ (i) \cong M_{k+i}$, $i = 0, 1$. Since

$$\begin{aligned}
sc_j s^*(t) &= \sin^2(\pi t/2)\cos(\pi t/2)\rho(u)(e_{1,1}^{(k)} \otimes e_{1,j}^{(k+1)})\rho(u)^*(t), \quad j = 1, 2, \ldots, k, \\
sc_j s^* c_i(t) &= \sin^2(\pi t/2)\cos(\pi t/2)\rho(u)(e_{1,i}^{(k)} \otimes e_{1,j}^{(k+1)})\rho(u)^*(t), \ \ i, j = 1, 2, \ldots, k, \\
sc_j(t) &= \sin(\pi t/2)\cos(\pi t/2)\rho(u)(t)(e_{1,j}^{(k)} \otimes e_{1,k+1}^{(k+1)}), \ \ j = 1, 2, \ldots, k,
\end{aligned}$$

for $t \in (0,1)$, we have that $C^*(\{c_j\}_j \cup \{s\})(t) = M_k \otimes M_{k+1}$ for $t \in (0,1)$.

Set $A = C^*(\{\tilde{c}_j\}_{j=1}^k \cup \{\tilde{s}\})$. Let $\Phi : A \to I(k, k+1)$ be the $*$-homomorphism defined by $\Phi(\tilde{c}_j) = c_j$ and $\Phi(\tilde{s}) = s$. It remains to show that Φ is injective. Let $(\pi, \mathcal{H})$ be an irreducible representation of A. Because for any $a \in A$ there exists an irreducible representation of A which preserves the norm of a (see [26, 4.3.10]), it suffices to show that there exists a representation φ of $I(k, k+1)$ on $\mathcal{H}$ such that $\varphi(c_j) = \pi(\tilde{c}_j)$ and $\varphi(s) = \pi(\tilde{s})$.

Set

$$\tilde{b} = \sum_{j=1}^{k} \tilde{c}_j^* \tilde{s}\tilde{s}^* \tilde{c}_j + \tilde{s}^* \tilde{s}.$$

By the following computations, we see that $\tilde{b}$ is in the center of A. Since $\{\tilde{c}_j\}_j \cup \{\tilde{s}\}$ satisfies the relations $\mathcal{R}_k$, in particular $\tilde{c}_1^2 = \tilde{c}_j \tilde{c}_j^*$, we have that

$$\begin{aligned}
\tilde{b}\tilde{c}_j &= \tilde{s}\tilde{s}^* \tilde{c}_j + \tilde{s}^* \tilde{s}\tilde{c}_j, \\
\tilde{c}_j \tilde{b} &= \tilde{s}\tilde{s}^* \tilde{c}_j + \tilde{c}_j \tilde{s}^* \tilde{s}, \\
\tilde{s}^* \tilde{s}\tilde{c}_j &= \tilde{c}_j - \tilde{c}_1^2 \tilde{c}_j = \tilde{c}_j \tilde{s}^* \tilde{s}, \\
\tilde{b}\tilde{s} &= \tilde{b}\tilde{c}_1 \tilde{s} = \tilde{s}\tilde{s}^* \tilde{s} + \tilde{s}^* \tilde{s}^2, \\
\tilde{s}\tilde{b} &= \sum \tilde{s}\tilde{c}_j^* \tilde{s}\tilde{s}^* \tilde{c}_j + \tilde{s}\tilde{s}^* \tilde{s}, \\
\tilde{s}^* \tilde{s}^2 &= \tilde{c}_1 \tilde{s} - \tilde{c}_1^3 \tilde{s} = \tilde{s} - \tilde{s} = 0, \\
\tilde{s}^* \tilde{s}\tilde{c}_j^* \tilde{s}\tilde{s}^* \tilde{c}_j &= \tilde{c}_j^* \tilde{s}\tilde{s}^* \tilde{c}_j - \tilde{c}_j^* \tilde{c}_j \tilde{c}_j^* \tilde{s}\tilde{s}^* \tilde{c}_j = 0.
\end{aligned}$$

Hence $[\tilde{b}, \tilde{c}_j] = 0$ and $[\tilde{b}, \tilde{s}] = 0$.

Set $\bar{c}_j = \pi(\tilde{c}_j)$, $\bar{s} = \pi(\tilde{s})$, and $\bar{b} = \pi(\tilde{b})$. Since $0 \leqq \tilde{b} \leqq 1$, we have that $\mathrm{sp}(\tilde{b}) = [0,1]$ and obtain $\beta \in [0,1]$ such that $\beta 1 = \bar{b}$. When $\beta = 0$ we have $\bar{s} = 0$. Thus $\{\bar{c}_j\}_j$ satisfies the relations for matrix units $\{e_{1,j}^{(k)}\}$ of M_k, and then

$\mathcal{H}=\mathbb{C}^k$. Set $V_0 : I(k,k+1)\to M_k$ as the irreducible representation at 0. Since $V_0(c_j)=e^{(k)}_{1,j}$ we obtain a unitary $u_0\in U(M_k)$ such that $\operatorname{Ad}u_0\circ V_0(c_j)=\bar{c}_j$ and define $\varphi=\operatorname{Ad}u_0\circ V_0$.

When $\beta=1$, by the following computations, we see that $\bar{c}_j^*\bar{c}_j$, $j=1,2,...,n$, and $\bar{s}^*\bar{s}$ are orthogonal projections. Since $\bar{b}=1$, we have that $\sum\bar{c}_j^*(1-\bar{s}\bar{s}^*)\bar{c}_j=0$. Then it follows that $\bar{c}_1^2=\bar{s}\bar{s}^*$, $\bar{c}_1^4=\bar{c}_1^2$, $(\bar{c}_j^*\bar{c}_j)^2=\bar{c}_j^*\bar{c}_1^2\bar{c}_j=\bar{c}_j^*\bar{c}_j$, and $(\bar{s}^*\bar{s})^2=\bar{s}^*\bar{c}_1^2\bar{s}=\bar{s}^*\bar{s}$. From $\sum\bar{c}_j^*\bar{c}_j+\bar{s}^*\bar{s}=1$ it follows that $\bar{c}_j^*\bar{c}_j$, $j=1,2,...,k$ and $\bar{s}^*\bar{s}$ are mutually orthogonal projections. Hence $\{\bar{c}_j\}_j\cup\{\bar{s}\}$ satisfies the relations for matrix units $\{e^{(k+1)}_{1,j}\}_{j=1}^{k+1}$ of M_{k+1}. Then we see that $\mathcal{H}=\mathbb{C}^{(k+1)}$ and can define $\varphi : I(k,k+1)\to B(\mathcal{H})$ as the irreducible representation of $I(k,k+1)$ at $t=1$ up to unitary equivalence.

When $0<\beta<1$, by the following computations, we see that

$$\begin{aligned}
E_{i,j} &= (\beta(1-\beta))^{-1}\bar{c}_i^*\bar{s}\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{c}_i,\\
E_{j,k+1} &= (\beta(1-\beta))^{-1}\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{s},
\end{aligned}$$

$i,j=1,2,...,k$, are mutually orthogonal projections. Since $\bar{b}\bar{s}=\bar{s}\bar{s}^*\bar{s}$ and $(1-\bar{b})\bar{s}^*\bar{s}=(1-\bar{s}^*\bar{s})\bar{s}^*\bar{s}$, we have that

$$\begin{aligned}
E_{i,j}^2 &= (\beta(1-\beta))^{-2}\bar{c}_i^*\bar{s}\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{c}_1^2\bar{s}\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{c}_i\\
&= (\beta(1-\beta))^{-2}\bar{c}_i^*\bar{s}\bar{s}^*\bar{s}\bar{c}_j^*\bar{c}_j\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{c}_i\\
&= \beta^{-1}(1-\beta)^{-2}\bar{c}_i^*\bar{s}(1-\bar{s}^*\bar{s})\bar{c}_j^*\bar{c}_j\bar{s}^*\bar{c}_i=E_{i,j},\\
E_{j,k+1}^2 &= (\beta(1-\beta))^{-2}(\bar{c}_j^*\bar{c}_j)^2(\bar{s}^*\bar{s})^2\\
&= \beta^{-1}(1-\beta)^{-2}\bar{c}_j^*\bar{c}_j(1-\bar{s}^*\bar{s})\bar{s}^*\bar{s}=E_{j,k+1},\\
\sum_{i,j}E_{i,j}+\sum_j E_{j,k+1} &= (\beta(1-\beta))^{-1}(\sum_{i=1}^k\bar{c}_i^*\bar{s}(1-\bar{s}^*\bar{s})\bar{s}^*\bar{c}_i+(1-\bar{s}^*\bar{s})\bar{s}^*\bar{s})\\
&= \beta^{-1}(\sum_{i=1}^k\bar{c}_i^*\bar{s}\bar{s}^*\bar{c}_i+\bar{s}^*\bar{s})=1.
\end{aligned}$$

Set

$$\begin{aligned}
&F_{i,j}=\beta^{-1}(1-\beta)^{-1/2}\bar{s}\bar{c}_j\bar{s}^*\bar{c}_i,\\
&F_{j,k+1}=(\beta(1-\beta))^{-1/2}\bar{s}\bar{c}_j,\quad i,j=1,2,...,k.
\end{aligned}$$

Then it follows that $F_{i,j}^*F_{i,j}=E_{i,j}$, $F_{j,k+1}^*F_{j,k+1}=E_{j,k+1}$, $F_{i,j}F_{i,j}^*=E_{1,1}$, and $F_{j,k+1}F_{j,k+1}^*=E_{1,1}$. Thus $\{F_{i,j}\}_{i,j}\cup\{F_{j,k+1}\}_j$ satisfies the same relations as matrix units $\{e^{(k)}_{1,i}\otimes e^{(k+1)}_{1,j}\}_{i,j}\cup\{e^{(k)}_{1,j}\otimes e^{(k+1)}_{1,k+1}\}_j$ of $M_k\otimes M_{k+1}$. It is not so hard to see that $\beta^{1/2}\sum_{j=1}^k F_{1,j}^*F_{j,k+1}=\bar{s}$, $\beta^{1/2}\sum_{j=1}^k F_{1,j}^*F_{i,j}=\bar{s}\bar{s}^*\bar{c}_i$, and $\beta\bar{c}_i=\bar{s}\bar{s}^*\bar{c}_i+\bar{s}^*\bar{s}\bar{c}_i$. Then we have that $C^*(\{F_{i,j}\}\cup\{F_{j,k+1}\})=C^*(\{\bar{c}_j\}_j\cup\{\bar{s}\})$ and $\mathcal{H}=\mathbb{C}^{k(k+1)}$. Let V_β be the irreducible representation of $I(k,k+1)$ at $t\in(0,1)$ with $\sin^2(\pi t/2)=\beta$. Then $V_\beta\circ\Phi(\tilde{b})=\beta$ and there exists a unitary u_β such that

$$\begin{aligned} F_{i,j} &= \beta^{-1}(1-\beta)^{-1/2}\operatorname{Ad}u_\beta \circ V_\beta(sc_j s^* c_i), \\ F_{j,k+1} &= (\beta(1-\beta))^{-1/2}\operatorname{Ad}u_\beta \circ V_\beta(sc_j). \end{aligned}$$

Hence, we have that $\operatorname{Ad}u_\beta \circ V_\beta(c_j) = \bar{c}_j$ and $\operatorname{Ad}u_\beta \circ V_\beta(s) = \bar{s}$ and obtain $\varphi = \operatorname{Ad}u_\beta \circ V_\beta$. This completes the proof. $\blacksquare$

Corollary 2.2. *Let A be a unital C^*-algebra. Suppose that A^1 contains $\bar{c}_i$, $i = 1, 2, ..., k$, and $\bar{s}$ satisfying the relations $\mathcal{R}_k$ and $\mathrm{sp}(|\bar{s}|) = [0,1]$. Then there exists a unital embedding $\varphi : I(k, k+1) \hookrightarrow A$ such that $\varphi(c_i) = \bar{c}_i$ and $\varphi(s) = \bar{s}$.*

Proof. By the universal property of $I(k, k+1)$, there exists a unital $*$ homomorphism $\varphi : I(k, k+1) \to A$ such that $\varphi(c_j) = \bar{c}_j$ and $\varphi(s) = \bar{s}$. We denote by X_φ the compact subset of $[0,1]$ determined by $\ker(\varphi) = \{f \in I(k, k+1); f|_{X_\varphi} = 0\}$. For any $f \in C([0,1])$ it follows that $f(|s|)(t) = f \circ \sin(\pi t/2)\ 1_k \otimes e_{k+1}^{(k+1)}$. Then we have that $f|_{\sin(\frac{\pi}{2}X_\varphi)} = 0 \Leftrightarrow f(|s|)|_{X_\varphi} = 0 \Leftrightarrow f(|\bar{s}|) = 0 \Leftrightarrow f|_{\mathrm{sp}(|\bar{s}|)} = 0$. Thus $\mathrm{sp}(|\bar{s}|) = \sin(\frac{\pi}{2}X_\varphi)$, which concludes that $\ker(\varphi) = 0$. $\blacksquare$

3 Property (SI) for projectionless C^*-algebras

Definition 3.1. Let A be a unital C^*-algebra and $\tau \in T(A)$. We recall the dimension function d_τ and define $\overline{d}_\tau : A^1_+ \to \mathbb{R}_+$ by

$$\begin{aligned} d_\tau(f) &= \lim_{n\to\infty} \tau((1/n + f)^{-1}f), \\ \overline{d}_\tau(f) &= \lim_{n\to\infty} \tau(f^n), \quad f \in A^1_+. \end{aligned}$$

Lemma 3.2. *For $f_n \in A^1_+$, $n \in \mathbb{N}$ with $(f_n)_n \in A_\infty$ and an increasing sequence m_n, $n \in \mathbb{N}$ with $m_n \nearrow \infty$, it follows that*

(1) *If $\lim\limits_{n\to\infty} \max\limits_{\tau\in T(A)} \tau(f_n) = 0$ then there exists $\tilde{f}_n \in A^1_+$, $n \in \mathbb{N}$ such that $(\tilde{f}_n)_n = (f_n)_n$ and $\lim\limits_{n\to\infty} \max\limits_{\tau\in T(A)} d_\tau(\tilde{f}_n) = 0$.*

(2) *There exists $\tilde{f}_n \in A^1_+$, $n \in \mathbb{N}$ such that $(\tilde{f}_n)_n = (f_n)_n$ and $\liminf\limits_{n\to\infty} \min\limits_{\tau\in T(A)} \overline{d}_\tau(\tilde{f}_n) \geq \liminf_n \min_\tau \tau(f_n^{m_n})$.*

(3) *If A absorbs $\mathcal{Z}$ tensorially, then there exists $f_n^{(i)} \in A^1_+$, $i = 0, 1$, $n \in \mathbb{N}$ such that $(f_n^{(i)})_n \in A_\infty$, $f_n^{(0)} f_n^{(1)} = 0$, $(f_n^{(i)})_n \leq (f_n)_n$, $i = 0, 1$, and $\liminf\limits_{n\to\infty} \min\limits_{\tau\in T(A)} \overline{d}_\tau(f_n^{(i)}) \geq \liminf_n \min_\tau \tau(f_n^{m_n})/2$.*

Proof. (1) Let $\varepsilon_n > 0$ be such that $\varepsilon_n \searrow 0$ and $\max_{\tau\in T(A)} \tau(f_n) \leq \varepsilon_n^2$. Set

$$g_\varepsilon(t) = \begin{cases} (1-\varepsilon)^{-1}(t-\varepsilon), & \varepsilon \leq t \leq 1, \\ 0, & 0 \leq t \leq \varepsilon, \end{cases}$$

and $\tilde{f}_n = g_{\varepsilon_n}(f_n)$. Then we have that $\|\tilde{f}_n - f_n\| \le \varepsilon_n$ and $\varepsilon_n \lim_{m\to\infty}(1/m + \tilde{f}_n)^{-1}\tilde{f}_n \le f_n$, which implies that $d_\tau(\tilde{f}_n) \le \varepsilon_n$, for any $\tau \in T(A)$.

(2) Let $\varepsilon_n > 0$ be such that $\varepsilon_n \searrow 0$, and $(1-\varepsilon_n)^{m_n} \to 0$. Set

$$g_\varepsilon(t) = \begin{cases} (1-\varepsilon)^{-1}t, & 0 \le t \le 1-\varepsilon, \\ 1, & 1-\varepsilon \le t \le 1, \end{cases}$$

and $\tilde{f}_n = g_{\varepsilon_n}(f_n)$. Then we have that $\|\tilde{f}_n - f_n\| \le \varepsilon_n$ and $f_n^{m_n} = f_n^{m_n}(\lim_{l\to\infty} \tilde{f}_n^l + \chi([0, 1-\varepsilon_n))(f_n)) \le \lim_{l\to\infty} \tilde{f}_n^l + (1-\varepsilon_n)^{m_n}$, (where $\chi(S)$ means the characteristic function of S,) which implies that $\tau(f_n^{m_n}) \le \overline{d}_\tau(\tilde{f}_n) + (1-\varepsilon_n)^{m_n}$, for any $\tau \in T(A)$.

(3) Set $c = \liminf_{n\to\infty} \min_{\tau\in T(A)} \tau(f_n^{m_n})$. Since $A \cong A \otimes \bigotimes_{n\in\mathbb{N}} \mathcal{Z}$, we obtain $l_n \in \mathbb{N}$ and $\overline{f}_n \in (A \otimes \bigotimes_{j=1}^{l_n} \mathcal{Z})^1_+$ such that $l_n \nearrow \infty$ and $m_n\|\overline{f}_n - f_n\| \to 0$, which implies that $(\overline{f}_n)_n \in A_\infty$ and $\liminf_{n\to\infty} \min_{\tau\in T(A)} \tau(\overline{f}_n^{m_n}) = c$. By an argument as in the proof of (2), we obtain $\tilde{f}_n \in (A \otimes \bigotimes_{l=1}^{l_n} \mathcal{Z})^1_+$, $n \in \mathbb{N}$ such that $(\tilde{f}_n)_n = (\overline{f}_n) = (f_n)$ and $\liminf_{n\to\infty} \min_{\tau\in T(A)} \overline{d}_\tau(\tilde{f}_n) \ge c$. Let $g_n^{(i)} \in \mathcal{Z}^1_+$, $i = 0, 1$, $n \in \mathbb{N}$ be such that $g_n^{(0)} g_n^{(1)} = 0$, $\liminf_n \overline{d}_{\tau_{\mathcal{Z}}}(g_n^{(i)}) = 1/2$, $i = 0, 1$, where $\tau_{\mathcal{Z}}$ means the unique tracial state of $\mathcal{Z}$. Set $f_n^{(i)} = \tilde{f}_n \otimes g_n^{(i)} \in A \otimes \bigotimes_{l=1}^{l_n+1} \mathcal{Z}$. Since $l_n \nearrow \infty$, it follows that $(f_n^{(i)})_n \in A_\infty$, and since $\tau((f_n^{(i)})^p) = \tau(\tilde{f}_n^p \otimes 1)\tau_{\mathcal{Z}}((g_n^{(i)})^p)$, $p \in \mathbb{N}$, $\tau \in T(A)$, it follows that $\liminf_n \min_\tau \overline{d}_\tau(f_n^{(i)}) = \liminf_n \min_\tau \overline{d}_\tau(\tilde{f}_n)\overline{d}_{\tau_{\mathcal{Z}}}(g_n^{(i)}) \ge c/2$, $i = 0, 1$. ∎

In [31], we have defined the technical condition, called property (SI), for C^*-algebra with non trivial projections. For projectionless C^*-algebras, we generalize this technical condition in the following.

Definition 3.3. We say that A has the *property (SI)* , when for any e_n and $f_n \in A^1_+$, $n \in \mathbb{N}$ satisfying the following conditions: $(e_n)_n, (f_n)_n \in A_\infty$,

$$\lim_{n\to\infty} \max_{\tau\in T(A)} \tau(e_n) = 0,$$

$$\liminf_{n\to\infty} \min_{\tau\in T(A)} \tau(f_n^n) > 0,$$

there exists $s_n \in A^1$, $n \in \mathbb{N}$, such that $(s_n)_n \in A_\infty$ and

$$(s_n^* s_n) = (e_n), \quad (f_n s_n) = (s_n).$$

Example 3.4. Any UHF algebra has the property (SI).

Proof. Let B be a UHF algebra, and let e_n and $f_n \in B^1_+$, $n \in \mathbb{N}$ satisfy the conditions in the property (SI). Let B_n, $n \in \mathbb{N}$ be an increasing sequence of matrix subalgebras of B such that $\overline{(\bigcup_{n\in\mathbb{N}} B_n)} = B$ and $1_{B_n} = 1_B$. For any B_n, we denote by $\Phi_n : B \longrightarrow B'_n \cap B$ the conditional expectations ([31, Section 1]). By $(e_n)_n$, $(f_n)_n \in B_\infty$, we obtain a slow increasing sequence $m_n \in \mathbb{N}$, $n \in \mathbb{N}$ such that $m_n \nearrow \infty$, $m_n \leq n$, $(\Phi_{m_n}(e_n))_n = (e_n)_n$, and

$$\lim_{n\to\infty} m_n \|\Phi_{m_n}(f_n) - f_n\| = 0,$$

and we obtain a fast increasing sequence l_n, $n \in \mathbb{N}$, and $\overline{e}_n, \overline{f}_n \in (B'_{m_n} \cap B_{l_n})^1_+$ such that $m_n < l_n$, $(\overline{e}_n)_n = (\Phi_{m_n}(e_n))_n$, and $\lim_{n\to\infty} m_n \|\overline{f}_n - \Phi_{m_n}(f_n)\| = 0$. Then we have that $\lim \tau(\overline{e}_n) = 0$ and $\lim \|\overline{f}_n^{m_n} - f_n^{m_n}\| = 0$, which implies that $\liminf \tau(\overline{f}_n^{m_n}) = \liminf \tau(f_n^{m_n}) > 0$. By Lemma 3.2 (1), we obtain $\tilde{e}_n \in (B'_{m_n} \cap B_{l_n})^1_+$ such that $(\tilde{e}_n)_n = (e_n)_n$ and $\lim d_\tau(\tilde{e}_n) = 0$. By Lemma 3.2 (2), we obtain $\tilde{f}_n \in (B'_{m_n} \cap B_{l_n})^1_+$ such that $(\tilde{f}_n)_n = (f_n)_n$ and

$$\liminf_{n\to\infty} \overline{d}_\tau(\tilde{f}_n) \geq \liminf \tau(\overline{f}_n^{m_n}) > 0.$$

Taking a large $N \in \mathbb{N}$, we have that

$$d_{\mathrm{Tr}_n}(\tilde{e}_n) = d_\tau(\tilde{e}_n) < \overline{d}_\tau(\tilde{f}_n) = \overline{d}_{\mathrm{Tr}_n}(\tilde{f}_n),\ n \geq N,$$

where Tr_n is the normalized trace of $B'_{m_n} \cap B_{l_n}$. Then, we obtain $s_n \in (B'_{m_n} \cap B_{l_n})^1$ such that $s_n^* s_n = \tilde{e}_n$, $\tilde{f}_n s_n = s_n$, hence we have that $(s_n)_n \in B_\infty$, $(s_n^* s_n)_n = (e_n)_n$, and $(f_n s_n)_n = (s_n)_n$. ■

By the above example and the following proposition, we see that the Jiang-Su algebra has the property (SI). This proposition is motivated by Lemma 3.3 in [23].

Proposition 3.5. *Let A be a unital C^*-algebra absorbing the Jiang-Su algebra $\mathcal{Z}$ tensorially. If $A \otimes B$ has the property (SI) for any UHF algebra B, then A also has the property (SI).*

In order to prove the above proposition, we define the projectionless C^*-algebra $\mathcal{Z}_k$ for $k \in \mathbb{N} \setminus \{1\}$ by

$$\begin{aligned}\mathcal{Z}_k = \quad & \{f \in C([0,1]) \otimes M_{k^\infty} \otimes M_{(k+1)^\infty}; \\ & f(0) \in M_{k^\infty} \otimes 1_{k+1^\infty},\ f(1) \in 1_{k^\infty} \otimes M_{k+1^\infty}\}.\end{aligned}$$

This projectionless C^*-algebra $\mathcal{Z}_k$ was introduced by Rørdam and Winter in [30], [34] as a mediator between C^*-algebras absorbing UHF algebras and C^*-algebras absorbing the Jiang-Su algebra.

Proof. Suppose that e_n and $f_n \in A^1_+$, $n \in \mathbb{N}$ satisfy the conditions in the property (SI). Let k be a natural number with $k \geq 2$, $B^{(i)}$ the UHF algebra of rank $(k+i)^\infty$, and $\Phi^{(i)}$ the canonical unital embeddings of $A \otimes B^{(i)}$ into $A \otimes B^{(0)} \otimes B^{(1)}$, $i = 0, 1$.

By Lemma 3.2 there exist $f_n^{(i)} \in A_+^1$, $i = 0, 1$, $n \in \mathbb{N}$, such that $(f_n^{(i)})_n \in A_\infty$, $(f_n^{(0)})_n (f_n^{(1)})_n = 0$, $(f_n^{(i)})_n \leq (f_n)_n$, and $\liminf_{n\to\infty} \min_{\tau \in T(A)} \tau({f_n^{(i)}}^n) \geq \liminf_n \min_\tau \overline{d}_\tau(f_n^{(i)}) \geq \liminf_n \min_\tau \tau(f_n^n)/2 > 0$, $i = 0, 1$.

Applying the property (SI) of $A \otimes B^{(i)}$ to $e_n \otimes 1_{(k+i)^\infty}$ and $f_n^{(i)} \otimes 1_{(k+i)^\infty} \in A \otimes {B^{(i)}}_+^1$ we obtain $s_n^{(i)} \in A \otimes {B^{(i)}}^1$, $i = 0, 1$, $n \in \mathbb{N}$ such that $(s_n^{(i)})_n \in (A \otimes B^{(i)})_\infty$,

$$({s_n^{(i)}}^* s_n^{(i)})_n = (e_n \otimes 1_{(k+i)^\infty}), \quad (f_n^{(i)} \otimes 1_{(k+i)^\infty} \cdot s_n^{(i)})_n = (s_n^{(i)}).$$

Note that $(\Phi^{(i)}({s_n^{(i)}}^* s_n^{(i)}))_n = (e_n \otimes 1_{k^\infty} \otimes 1_{(k+1)^\infty})_n$, $(f_n \otimes 1_{k^\infty} \otimes 1_{(k+1)^\infty})_n \cdot (\Phi^{(i)}(s_n^{(i)}))_n = (\Phi^{(i)}(s_n^{(i)}))_n$, and $(\Phi^{(0)}(s_n^{(0)}))_n^* \ (\Phi^{(1)}(s_n^{(1)}))_n = 0$.

Define $s_n \in A \otimes {\mathcal{Z}_k}^1$, $n \in \mathbb{N}$ by

$$s_n(t) = \cos(\pi t/2)\Phi^{(0)}(s_n^{(0)}) + \sin(\pi t/2)\Phi^{(1)}(s_n^{(1)}), \quad t \in [0, 1].$$

Since

$$\begin{array}{rcl} (s_n^* s_n(t))_n &=& (\cos^2(\pi t/2)\Phi^{(0)}({s_n^{(0)}}^* s_n^{(0)}) + \sin^2(\pi t/2)\Phi^{(1)}({s_n^{(1)}}^* s_n^{(1)}) \\ &+& \cos \cdot \sin(\pi t/2)(\Phi^{(0)}(s_n^{(0)})^* \Phi^{(1)}(s_n^{(1)}) + \Phi^{(1)}(s_n^{(1)})^* \Phi^{(0)}(s_n^{(0)})))_n \\ &=& (e_n \otimes 1_{k^\infty} \otimes 1_{(k+1)^\infty})_n, \quad t \in [0, 1], \end{array}$$

$(f_n \otimes 1_{k^\infty} \otimes 1_{(k+1)^\infty})_n \ (s_n(t)) = (s_n(t))$, $t \in [0, 1]$, and $\mathrm{Lip}(s_n) = \pi$, $n \in \mathbb{N}$, it follows that

$$(s_n^* s_n)_n = (e_n \otimes 1_{\mathcal{Z}_k})_n, \quad (f_n \otimes 1_{\mathcal{Z}_k})_n (s_n)_n = (s_n)_n.$$

Set $\iota : A_\infty \hookrightarrow (A \otimes \mathcal{Z}_k)_\infty$ by $\iota((a_n)_n) = (a_n \otimes 1_{\mathcal{Z}_k})_n$. Since $A \cong A \otimes \bigotimes_{n\in\mathbb{N}} \mathcal{Z}$ and $\mathcal{Z}_k \subset_{\mathrm{unital}} \mathcal{Z}$, for any finite subset $F \subset A_\infty$, we obtain a unital embedding $\Phi_F : (A \otimes \mathcal{Z}_k)_\infty \hookrightarrow A_\infty$ such that $\Phi_F \circ \iota(x) = x$, $x \in F$.

Define $s = \Phi_{\{(e_n),(f_n)\}}((s_n)_n) \in A_\infty$, then we conclude that $s^* s = (e_n)$ and $(f_n)s = s$. ∎

4 $\mathcal{Z}$ absorption of Crossed products.

In this section we prove Theorem 1.2. We denote by A_α the fixed point algebra of $\alpha \in \mathrm{Aut}(A)$ and by α_∞ the automorphism of A_∞ induced by α. In the following Lemma 4.1, mimicking Theorem 4.4 in [12], we use the weak Rohlin property to obtain a set of elements in $(A_\infty)_{\alpha_\infty}$ which satisfies the same relations as $\{c_j\}_{j=1}^k$ in $\mathcal{R}_k$. After that, applying the property (SI) and the weak Rohlin property, we obtain the generators of prime dimension drop algebras in $(A_\infty)_{\alpha_\infty}$.

Lemma 4.1. *Let A be a unital separable simple C^*-algebra which has a unique tracial state τ and absorbs the Jiang-Su algebra tensorially. Suppose that $\alpha \in$*

$\mathrm{Aut}(A)$ *has the weak Rohlin property. Then for any* $k \in \mathbb{N}$ *there exist* $c_{j,n} \in A$, $j = 1, 2, ..., k$, $n \in \mathbb{N}$ *such that* $(c_{j,n})_n \in (A_\infty)^1_{\alpha_\infty}$,

$$(c_{1,n})_n \geqq 0, \quad (c_{i,n})_n (c_{j,n})^* = \delta_{i,j}(c_{1,n})^2,$$

$$\|(c_{1,n})_n\| = 1, \quad \|1 - \sum_{j=1}^{k} (c_{j,n})^*_n (c_{j,n})\| = 1, \quad \lim_{n\to\infty} \tau(c^n_{1,n}) = 1/k.$$

(which implies $\lim_{n\to\infty} \tau(1_A - \sum_{j=1}^{k} c^*_{j,n} c_{j,n}) = 0$.*)*

Proof. Let Φ_m, $m \in \mathbb{N}$ be the unital embeddings of $\mathcal{Z}$ into $A \otimes \bigotimes_{m\in\mathbb{N}} \mathcal{Z} \cong A$ defined by

$$\Phi_m(x) = 1_A \otimes 1_{\bigotimes_{i=1}^{m} \mathcal{Z}} \otimes x \otimes 1_{\bigotimes_{i=m+2}^{\infty} \mathcal{Z}}, \quad x \in \mathcal{Z},$$

and Φ be the unital embedding of $\mathcal{Z}$ into A_∞ defined by $\Phi(x) = (\Phi_m(x))_m$, $x \in \mathcal{Z}$. Remark that $\tau(\Phi_m(x)) = \tau_{\mathcal{Z}}(x)$, $m \in \mathbb{N}$, $x \in \mathcal{Z}$.

In the definition of $c_j \in I(k, k+1)$ in Section 2, replacing cos with $\cos^{1/n^2}$ we obtain $c^{(j)}_n \in I(k, k+1) \subset \mathcal{Z}$, $j = 1, 2, ..., k$ such that

$$c^{(1)}_n \geqq 0, \quad c^{(i)}_n {c^{(j)}_n}^* = \delta_{i,j} {c^{(1)}_n}^2,$$

$$\|(c^{(1)}_n)_n\| = 1, \quad \|1 - \sum_{j=1}^{k} {c^{(j)}_n}^* c^{(j)}_n\| = 1, \quad \tau_{\mathcal{Z}}({c^{(1)}_n}^n) \nearrow 1/k.$$

Let $\varepsilon_n > 0$, $n \in \mathbb{N}$ be such that $\varepsilon_n \searrow 0$ and $\tau_{\mathcal{Z}}({c^{(1)}_n}^n) > 1/k - \varepsilon_n$, $n \in \mathbb{N}$.

Let k_n and $l_n \in \mathbb{N}$, $n \in \mathbb{N}$ be such that $l_n \nearrow \infty$ and $l^2_n < k_n$. Since $\alpha \in \mathrm{Aut}(A)$ has the weak Rohlin property and A and $\mathcal{Z}$ are separable C^*-algebras, we obtain $f^{(n)} = (f^{(n)}_m)_m \in (A \cup \bigcup_{j\in\mathbb{Z}} \alpha^j_\infty(\Phi(\mathcal{Z})))' \cap A^\infty$ such that $\|f^{(n)}\| = 1$, and

$$\begin{aligned} &\alpha^p_\infty(f^{(n)}) f^{(n)} = 0, \quad 0 < |p| \leq 2(k_n + l_n), \\ &\tau({f^{(n)}_r}^n) > 1/(2(k_n + l_n) + 1) - \varepsilon_m, \quad r \geq m. \end{aligned}$$

Note that any subsequence of $(f^{(n)}_m)_m$ satisfies the above conditions. Then, taking a subsequence of $(f^{(n)}_m)_m$, we may suppose that

$$|\tau(\Phi_m({c^{(1)}_n}^n) \cdot {f^{(n)}_m}^n) - \tau_{\mathcal{Z}}({c^{(1)}_n}^n) \tau({f^{(n)}_m}^n)| < \varepsilon_m.$$

For $p \in \mathbb{Z}$, define $a_{p,n} \geq 0$ by

$$a_{p,n} = \begin{cases} 1 - (|p| - k_n)/l_n, & k_n < |p| \leq k_n + l_n, \\ 1, & |p| \leq k_n, \\ 0, & k_n + l_n < |p|, \end{cases}$$

and define completely positive maps $\varphi_n : \mathcal{Z} \longrightarrow A_\infty$, by

$$\varphi_n(x) = \sum_{|p|\leq k_n+l_n} a_{p,n}\alpha^p_\infty(\Phi(x))\alpha^p_\infty(f^{(n)}).$$

Then we have that $\|\alpha_\infty(\varphi_n(x)) - \varphi_n(x)\|$

$$\begin{aligned} &= \| \sum_{|p|\leq k_n+l_n} (a_{p,n} - a_{p-1,n}) \cdot \alpha^p_\infty(\Phi(x))\alpha^p_\infty(f^{(n)})\| \\ &= \|x\|/l_n, \quad x \in \mathcal{Z}, \end{aligned}$$

$$\begin{aligned} \varphi_n(c^{(i)}_n)\varphi_n(c^{(j)}_n)^* &= \sum_{|p|\leq k_n+l_n} a^2_{p,n}\alpha^p_\infty(\Phi(c^{(i)}_n {c^{(j)}_n}^*))\alpha^p_\infty(f^{(n)})^2 \\ &= \delta_{i,j}\varphi_n(c^{(1)}_n)^2, \quad n \in \mathbb{N}, \end{aligned}$$

$\|\varphi_n(c^{(1)}_n)\| = 1$, and $\|1 - \sum_{j=1}^k \varphi_n(c^{(j)}_n)^*\varphi_n(c^{(j)}_n)\ \| = 1$.

Let $c^{(j)}_{n,m} \in A^1$, $j = 1,2,...,k$, $m \in \mathbb{N}$ be components of $\varphi_n(c^{(j)}_n)$ (i.e., $(c^{(j)}_{n,m})_m = \varphi_n(c^{(j)}_n) \in A_\infty$) with $c^{(1)}_{n,m} \geqq 0$, then we have that

$$\begin{aligned} \liminf_{m\to\infty} \tau({c^{(1)}_{n,m}}^n) &= \liminf_m \sum_{|p|\leq k_n+l_n} a^n_{p,n}\tau(\alpha^p(\Phi_m(c^{(1)}_n)^n)\alpha^p(f^{(n)}_m)^n) \\ &> (2k_n+1)\liminf_m \tau(\Phi_m({c^{(1)}_n}^n)\cdot {f^{(n)}_m}^n) \\ &= (2k_n+1)\liminf_m \tau_{\mathcal{Z}}({c^{(1)}_n}^n)\tau({f^{(n)}_m}^n) \\ &> \frac{(2k_n+1)}{(2(k_n+l_n)+1)}(1/k-\varepsilon_n) \to_{n\to\infty} 1/k, \end{aligned}$$

Let F_n be an increasing sequence of finite subsets of A^1 with $\overline{\bigcup_n F_n} = A^1$. By the above conditions, we obtain an increasing sequence $m_n \in \mathbb{N}$, $n \in \mathbb{N}$ such that

$$\begin{aligned} &\|[c^{(j)}_{n,m_n}, x]\| < \varepsilon, \quad x \in F_n, \\ &\|\alpha(c^{(j)}_{n,m_n}) - c^{(j)}_{n,m_n}\| < 1/l_n + \varepsilon_n, \\ &\|c^{(i)}_{n,m_n}{c^{(j)}_{n,m_n}}^* - \delta_{i,j}{c^{(1)}_{n,m_n}}^2\| < \varepsilon_n,\ i,j = 1,2,...,k, \\ &\|c^{(1)}_{n,m_n}\| > 1-\varepsilon_n, \quad \|1 - \sum_{j=1}^k {c^{(j)}_{n,m_n}}^* c^{(j)}_{n,m_n}\| > 1-\varepsilon_n, \\ &\tau({c^{(1)}_{n,m_n}}^n) > \frac{2k_n+1}{2(k_n+l_n)+1}(1/k-\varepsilon_n). \end{aligned}$$

Define $c_{j,n} = c^{(j)}_{n,m_n}$, $j = 1,2\ ,...,k$, then we have that $(c_{j,n})_n \in (A_\infty)_{\alpha_\infty}$,

$$(c_{1,n})_n \geqq 0, \quad (c_{i,n})_n(c_{j,n})^* = \delta_{i,j}(c_{1,n})^2,$$

$$\|(c_{1,n})_n\| = 1, \quad \|1 - \sum_{j=1}^{k} (c_{j,n})_n^*(c_{j,n})\| = 1, \quad \lim_{n\to\infty} \tau(c_{1,n}^n) = 1/k.$$

∎

By the technique in the proof of Lemma 4.6 [21] we obtain a generator s in $(A_\infty)_{\alpha_\infty}$ satisfying the relations in $\mathcal{R}_k$ together with $\{(c_{j,n})_n\}$ above. In the proof of the following proposition, $x \approx_\varepsilon y$ means $\|x - y\| < \varepsilon$.

Proposition 4.2. *Let A be a unital C^*-algebra which has a unique tracial state τ, absorbs the Jiang-Su algebra $\mathcal{Z}$ tensorially, and has the property (SI). Suppose that $\alpha \in \mathrm{Aut}(A)$ has the weak Rohlin property. Then for any $k \in \mathbb{N}$ there exists a set of norm-one elements $\{c_j\}_{j=1}^k \cup \{s\}$ in $(A_\infty)_{\alpha_\infty}$ satisfying $\mathcal{R}_k$.*

Proof. By Lemma 4.1 we obtain $c_m^{(j)} \in A^1$, $j = 1, 2, ..., k$, $m \in \mathbb{N}$ such that $(c_m^{(j)})_m \in (A_\infty)_{\alpha_\infty}$, $(c_m^{(1)})_m \geq 0$, $(c_m^{(i)})_m(c_m^{(j)})^* = \delta_{i,j}(c_m^{(1)})^2$, $\|(c_m^{(j)})_m\| = 1$, $\|1 - \sum_{j=1}^k (c_m^{(j)})^* \, (c_m^{(j)})\| = 1$, $\lim_{m\to\infty} \tau({c_m^{(1)}}^m) = 1/k$, and $\lim_{m\to\infty} \tau(1 - \sum {c_m^{(j)}}^* c_m^{(j)}) = 0$, where τ is a unique tracial state of A. Let $\varepsilon_m > 0$, $m \in \mathbb{N}$ be such that $\varepsilon_m \searrow 0$ and $\tau({c_m^{(1)}}^m) \geq 1/k - \varepsilon_m$.

Because of the weak Rohlin property of $\alpha \in \mathrm{Aut}(A)$ we obtain $f_m^{(l)} \in A_+^1$, $l, m \in \mathbb{N}$, such that $(f_m^{(l)})_m \in A_\infty$ and

$$\begin{aligned}
&\alpha_\infty^p((f_m^{(l)})_m)(f_m^{(l)}) = 0, \ p = 1, 2, ..., l-1, \\
&\|[f_r^{(l)}, c_m^{(1)}]\| < \varepsilon_m, \ r \geq m, \\
&\tau({f_r^{(l)}}^m) > 1/l - \varepsilon_m, \ r \geq m.
\end{aligned}$$

Note that any subsequence of $(f_m^{(l)})_m$ satisfies the above conditions. Since τ is the unique tracial state, taking a subsequence of $(f_m^{(l)})_m$ we may suppose that $\tau((c_m^{(1)} f_m^{(l)})^m) \approx_{\varepsilon_m} \tau({c_m^{(1)}}^m {f_m^{(l)}}^m) \approx_{\varepsilon_m} \tau({c_m^{(1)}}^m)\tau({f_m^{(l)}}^m) \geq 1/(kl) - 2\varepsilon_m$. Set

$$g_m^{(l)} = {c_m^{(1)}}^{1/2} f_m^{(l)} {c_m^{(1)}}^{1/2} \in A_+^1, \quad m \in \mathbb{N},$$

then we have that $(g_m^{(l)})_m \in {A_\infty}_+^1$, $l \in \mathbb{N}$ and $\liminf_{m\to\infty} \tau({g_m^{(l)}}^m) \geq 1/(kl)$.

By the property (SI) of A, we obtain ${s'}_m^{(l)} \in A^1$, $m \in \mathbb{N}$, such that $({s'}_m^{(l)})_m \in A_\infty$, $({{s'}_m^{(l)}}^* {s'}_m^{(l)}) = (1 - \sum_{j=1}^k {c_m^{(j)}}^* c_m^{(j)})$, and $(g_m^{(l)} {s'}_m^{(l)}) = ({s'}_m^{(l)})$. Remark that $(g_m^{(l)}) = (f_m^{(l)})(c_m^{(1)}) \leq (f_m^{(l)})$, $(g_m^{(l)}) \leq (c_m^{(1)})$, $(f_m^{(l)})({s'}_m^{(l)}) = ({s'}_m^{(l)})$, and $(c_m^{(1)})_m({s'}_m^{(l)}) = ({s'}_m^{(l)})$. Let $L_n \in \mathbb{N}$ be such that $2L_n^{-1/2} < \varepsilon_n$ ($L_n \nearrow \infty$) and define

$$s_m^{(L_n)} = L_n^{-1/2} \sum_{p=0}^{L_n - 1} \alpha^p({s'}_m^{(L_n)}).$$

Then we have that

$$\|\alpha(s_m^{(L_n)}) - s_m^{(L_n)}\| \leq 2L_n^{-1/2} < \varepsilon_n, \quad m \in \mathbb{N}.$$

Let $m_n \in \mathbb{N}$, $n \in \mathbb{N}$ be an increasing sequence with $m_n \nearrow \infty$ such that $(s_{m_n}^{(L_n)})_n \in A_\infty$, $\|s_{m_n}^{(L_n)}\| \leq 1 + \varepsilon_n$, $\|f_{m_n}^{(L_n)} {s'}_{m_n}^{(L_n)} - {s'}_{m_n}^{(L_n)}\| < \varepsilon_n / L_n$, $\|\alpha^p(f_{m_n}^{(L_n)})\ f_{m_n}^{(L_n)}\| < \varepsilon_n / L_n$, $\|{{s'}_{m_n}^{(L_n)}}^* {s'}_{m_n}^{(L_n)} - (1 - \sum_{j=1}^k {c_{m_n}^{(j)}}^* c_{m_n}^{(j)})\| \leq \varepsilon_n$, $\|\alpha^p(c_{m_n}^{(j)}) - c_{m_n}^{(j)}\| < \varepsilon_n/(2k)$, $j = 1, 2, ..., k$, $p = 1, 2, ..., L_n - 1$, and $\|c_{m_n}^{(1)} s_{m_n}^{(L_n)} - s_{m_n}^{(L_n)}\| < \varepsilon_n / L_n$, and set $s_n = s_{m_n}^{(L_n)}$.

Then we have that $\alpha_\infty((s_n)_n) = (s_n) \in A_\infty$,

$$\begin{aligned}
s_n^* s_n &\approx_{2\varepsilon_n} L_n^{-1} (\sum_{p=0}^{L_n - 1} \alpha^p({{s'}_{m_n}^{(L_n)}}^* f_{m_n}^{(L_n)}))(\sum_{q=0}^{L_n - 1} \alpha^q(f_{m_n}^{(L_n)} {s'}_{m_n}^{(L_n)})) \\
&\approx_{\varepsilon_n} L_n^{-1} \sum_{p=0}^{L_n - 1} \alpha^p({{s'}_{m_n}^{(L_n)}}^* {f_{m_n}^{(L_n)}}^2 {s'}_{m_n}^{(L_n)}) \\
&\approx_{\varepsilon_n} L_n^{-1} \sum \alpha^p({{s'}_{m_n}^{(L_n)}}^* {s'}_{m_n}^{(L_n)}) \\
&\approx_{\varepsilon_n} L_n^{-1} \sum \alpha^p(1 - \sum_{j=1}^k {c_{m_n}^{(j)}}^* c_{m_n}^{(j)}) \\
&\approx_{\varepsilon_n} 1 - \sum_{j=1}^k {c_{m_n}^{(j)}}^* c_{m_n}^{(j)},
\end{aligned}$$

$\|(s_n)_n\| = 1$, and $(c_{m_n}^{(1)})(s_n) = (s_n)$. Hence we conclude that $\{(c_{m_n}^{(j)})_n\}_{j=1}^k \cup \{(s_n)_n\} \subset (A_\infty)_{\alpha_\infty}^1$ and they satisfy the relations $\mathcal{R}_k$. ∎

Proof of Theorem 1.2 Applying Proposition 2.2 in [33] to $(A \times_\alpha \mathbb{Z})_\infty$ it suffices to show the following Lemma.

Lemma 4.3. *Let A be a unital separable projectionless C^*-algebra which has a unique tracial state and absorbs the Jiang-Su algebra tensorially. Suppose that A has property (SI) and $\alpha \in \mathrm{Aut}(A)$ has the weak Rohlin property. Then for any $k \in \mathbb{N}$ there exists a unital embedding Φ_k of $I(k, k+1)$ into $(A_\infty)_{\alpha_\infty}$.*

Proof. By Proposition 4.2 we obtain a set of norm one generators $\{c_j\}_{j=1}^k \cup \{s\}$ in $(A_\infty)_{\alpha_\infty}$ satisfying $\mathcal{R}_k$. Since $|s|$ and c_1 are norm one elements we have that $\mathrm{sp}(|s|) \supset \{0, 1\}$, and since A_∞ is a projectionless C^*-algebra we have that $\mathrm{sp}(|s|) = [0, 1]$. Then, by Corollary 2.2, we conclude the above statement. ∎

5 Stability for automorphisms

In this section, using the weak Rohlin property, we show the stability for automorphisms of the Jiang-Su algebra, Theorem 5.3.

First, we recall the generalized determinant introduced by P. de la Harpe and G. Skandalis (see [11], [19], [27]). Let A be a unital C^*-algebra with a unique tracial state τ. For any piecewise differentiable path $\xi : [0, 1] \to U(A)$, we set

$$\widetilde{\Delta}_\tau(\xi) = \frac{1}{2\pi\sqrt{-1}} \int_0^1 \tau(\dot{\xi}(t)\xi^*(t))dt \in \mathbb{R}.$$

When $\xi(0) = \xi(1) = 1$ we have that $\widetilde{\Delta}_\tau(\xi) \in \tau(K_0(A))$. For any $u \in U_0(A)$, there exists a piecewise differentiable path $\xi_u : [0,1] \to U(A)$ such that $\xi_u(0) = 1$, $\xi_u(1) = u$. The generalized determinant Δ_τ associated with the tracial state τ is the map from $U_0(A)$ to $\mathbb{R}/\tau(K_0(A))$ defined by $\Delta_\tau(u) = \widetilde{\Delta}(\xi_u) + \tau(K_0(A))$. Remark that Δ_τ is a group homomorphism.

Mimicking the proof of Lemma 6.2 in [19] we prove the following proposition. Hereinafter, we let log be the standard branch defined on the complement of the negative real axis.

Proposition 5.1. *Let B be the UHF algebra of rank k^∞, where $k \in \mathbb{N} \setminus \{1\}$, τ the unique tracial state of B, $\beta \in \mathrm{Aut}(B)$, and $u_n \in U(B)$, $n \in \mathbb{N}$. Suppose that $\beta \in \mathrm{Aut}(B)$ has the Rohlin property and*

$$\Delta_\tau(u_n) = 0, \quad \textit{for any } n \in \mathbb{N}.$$

Then there exist $v_n \in U(B)$, $n \in \mathbb{N}$ such that $(v_n)_n \in B_\infty$,

$$(v_n\beta(v_n)^*)_n = (u_n)_n, \quad \tau \circ \log(v_n\beta(v_n)^*u_n^*) = 0, \ \textit{for any } n \in \mathbb{N}.$$

The following lemma was essentially proved in [11].

Lemma 5.2. *Let A be a unital C^*-algebra with a unique tracial state τ.*

(1) For u_1, $u_2 \in U(A)$ with, $\|u_i - 1\| < 1/2$, $i = 1, 2$ it follows that

$$\tau \circ \log(u_1u_2) = \tau \circ \log(u_1) + \tau \circ \log(u_2).$$

(2) For u_1, u_2, and $v \in U(A)$ with $\|u_1 - u_2\| < 1/2$ and $\|v - 1\| < 1/4$, it follows that

$$\tau \circ \log(u_1vu_2^*v^*) = \tau \circ \log(u_1u_2^*).$$

Proof. (1) Let $h_i \in A_{\mathrm{sa}}$ be such that $\exp(2\pi\sqrt{-1}h_i) = u_i$, $i = 1, 2$, and $h_3 \in A_{\mathrm{sa}}$ be such that $\exp(2\pi\sqrt{-1}h_3) = u_1u_2$. Set $u(t) = \exp(2\pi\sqrt{-1}th_1) \cdot \exp(2\pi\sqrt{-1}th_2)$, $w(t) = \exp(2\pi\sqrt{-1}th_3)$, $t \in [0,1]$. Since $\|1 - u(t)\| < 1$, $\|1 - w(t)\| < 1$, and $\|1 - w^*u(t)\| < 2$, $t \in [0,1]$, we can define $h \in C([0,1]) \otimes A_{\mathrm{sa}}$ by $h(t) = \log(w^*u(t))$, $t \in [0,1]$, then u and w are homotopic, by $H(s,t) = w(t)\exp((1-s)h(t))$ with fixed endpoints $H(s,0) = 1$ and $H(s,1) = w(1)$. Hence, we have that

$$\begin{aligned}\tau \circ \log(u_1u_2) &= 2\pi\sqrt{-1}\tau(h_3) = \int_0^1 \tau(\dot{w}w^*(t))dt = \int_0^1 \tau(\dot{u}u^*(t))dt \\ &= 2\pi\sqrt{-1}\tau(h_1 + h_2) = \tau \circ \log(u_1) + \tau \circ \log(u_2).\end{aligned}$$

(2) Set $U_1 = v^*u_1vu_1^*$, $U_2 = u_1u_2^*$, then it follows that $\|U_i - 1\| < 1/2$, $i = 1, 2$. Applying (1), since $\tau \circ \log(U_1) = \tau \circ \log(v^*) + \tau \circ \log(u_1vu_1^*) = 0$ we have that $\tau \circ \log(U_1U_2) = \tau \circ \log(U_1) + \tau \circ \log(U_2) = \tau \circ \log(U_2)$. ∎

proof of Proposition 5.1. Because $\beta \in \mathrm{Aut}(B)$ has the Rohlin property in [20], there exist ${v'}_n \in U(B)$, $n \in \mathbb{N}$ such that $({v'}_n)_n \in B_\infty$, and

$$({v'}_n \beta({v'}_n)^*)_n = (u_n).$$

By the assumption and

$$\begin{aligned}
&\frac{1}{2\pi\sqrt{-1}}\tau \circ \log({v'}_n \beta({v'}_n)^* u_n^*) + \tau(K_0(B)) \\
&= \Delta_\tau(v'_n\beta(v'_n)^* u_n^*) = \Delta_\tau(v'_n\beta(v'_n)^*) - \Delta_\tau(u_n) = -\Delta_\tau(u_n),
\end{aligned}$$

we have that

$$\frac{1}{2\pi\sqrt{-1}}\tau \circ \log({v'}_n \beta({v'}_n)^* u_n^*) \in \tau(K_0(B)),\ n \in \mathbb{N}.$$

Since B is the UHF algebra of rank k^∞, we obtain $l_n \in \mathbb{N}$ and $m_n \in \mathbb{Z}$ such that $(m_n, k) = 1$ and

$$k^{-l_n} m_n = -\frac{1}{2\pi\sqrt{-1}}\tau \circ \log({v'}_n \beta({v'}_n)^* u_n^*) \in \tau(K_0(B)).$$

Set $\lambda_n = \exp(2\pi\sqrt{-1}k^{-l_n}m_n)$, then we have that $\lambda_n \to 1$, by $({v'}_n\ \beta({v'}_n)^*\ u_n^*)_n = 1$. By the Rohlin property of $\beta \in \mathrm{Aut}(B)$, there exist $p_n \in P(B)$ and $z_n \in U(B)$, $n \in \mathbb{N}$ such that $(p_n)_n \in B_\infty$, $(z_n)_n = 1_{B_\infty}$, and

$$\sum_{j=0}^{k^{l_n}-1} (\mathrm{Ad}\, z_n \circ \beta)^j (p_n) = 1_B.$$

Define

$$\begin{aligned}
\overline{v}_n &= \sum_{j=0}^{k^{l_n}-1} \exp(2\pi\sqrt{-1}jk^{-l_n}m_n) \cdot (\mathrm{Ad}\, z_n \circ \beta)^j (p_n), \\
v_n &= {v'}_n \overline{v}_n \in U(B),\ n \in \mathbb{N}.
\end{aligned}$$

Taking a subsequence of $(p_n)_n$, we may suppose that $(\overline{v}_n)_n \in B_\infty$. Then it follows that $(v_n)_n \in B_\infty$. By the definition of $\overline{v}_n$ we have that $\overline{v}_n \,\mathrm{Ad}\, z_n \circ \beta(\overline{v}_n)^*$ $= \lambda_n$ and

$$(v_n\beta(v_n)^* u_n^*)_n = (v_n \,\mathrm{Ad}\, z_n \circ \beta(v_n^*) u_n^*)_n = (\lambda_n {v'}_n \beta({v'}_n)^* u_n^*)_n = 1.$$

And, by Lemma 5.2, we have that $\tau \circ \log(v_n\beta(v_n)^* u_n^*)$

$$\begin{aligned}
&= \tau \circ \log(v_n \,\mathrm{Ad}\, z_n \circ \beta(v_n)^* u_n^*) \\
&= \tau \circ \log(\overline{v}_n \,\mathrm{Ad}\, z_n \circ \beta(\overline{v}_n)^* \,\mathrm{Ad}\, z_n \circ \beta({v'}_n)^* u_n^* {v'}_n) \\
&= 2\pi\sqrt{-1}k^{-l_n}m_n + \tau \circ \log({v'}_n \beta({v'}_n)^* u_n^*) = 0,\ n \in \mathbb{N}.
\end{aligned}$$

∎

Theorem 5.3. *Suppose that* $\sigma \in \mathrm{Aut}(\mathcal{Z})$ *has the weak Rohlin property and* $u_n \in U(\mathcal{Z})$, $n \in \mathbb{N}$ *satisfy* $(u_n)_n \in \mathcal{Z}_\infty$ *and that*

$$\Delta_{\tau_{\mathcal{Z}}}(u_n) = 0, \text{ for any } n \in \mathbb{N}.$$

Then there exist $v_n \in U(\mathcal{Z})$, $n \in \mathbb{N}$ *such that* $(v_n)_n \in \mathcal{Z}_\infty$ *and*

$$(v_n \sigma(v_n)^*)_n = (u_n)_n.$$

The following lemma is a direct adaptation of Proposition 4.6 in [19].

Lemma 5.4. *For any* $c > 0$ *there exists* $c' > 0$ *such that the following holds. Let* B *be a UHF algebra,* τ *the unique tracial state of* B*. Suppose that* $\widetilde{u}_n \in U(C([0,1]) \otimes B)$, $n \in \mathbb{N}$ *satisfies that* $(\widetilde{u}_n)_n \in (C([0,1]) \otimes B)_\infty$, $(\widetilde{u}_n(i))_n = 1$, $i = 0, 1$,

$$\widetilde{\Delta}_\tau(\widetilde{u}_n) = 0, \quad \tau \circ \log(\widetilde{u}_n(i)) = 0, \ i = 0,1, \ n \in \mathbb{N},$$

and $\mathrm{Lip}(\widetilde{u}_n) < c$, $n \in \mathbb{N}$*. Then there exist* $y_n \in U(C([0,1]^2) \otimes B)$, $n \in \mathbb{N}$ *such that* $(y_n)_n \in (C([0,1]^2) \otimes B)_\infty$,

$$y_n(0,t) = 1_B, \quad y_n(1,t) = \widetilde{u}_n(t), \ t \in [0,1],$$

$$y_n(s,i) = \exp(\log(\widetilde{u}_n(i))s), \ i = 0,1, \ s \in [0,1],$$

and $\mathrm{Lip}(y_n) < c'$, $n \in \mathbb{N}$.

Proof. Set $\partial E = \{(s,t) \in [0,1]^2; \ \{s,t\} \cap \{0,1\} \neq \phi\}$. By Proposition 4.6 in [19], for $c > 0$, we obtain $c' > 0$ satisfying that: for any AF-algebra A and for any $z \in U(C(\partial E) \otimes A)$ with $z(0,0) = 1$, $\mathrm{Lip}(z) < c$, and $[z]_1 = 0 \in K_1(C(\partial(E)) \otimes A)$, there exist $\widetilde{z} \in U(C([0,1]^2) \otimes A)$ such that $\widetilde{z}|_{\partial E} = z$ and $\mathrm{Lip}(\widetilde{z}) < c'$. Suppose that $\widetilde{u}_n \in U(C([0,1]) \otimes B)$ satisfies the conditions in the lemma. Define $U_n \in U(C(\partial E) \otimes B)$ by

$$U_n(s,t) = \begin{cases} 1, & s = 0, \\ \widetilde{u}_n(t), & s = 1, \\ \exp(\log(\widetilde{u}_n(i))s), & t = i, \ i = 0, 1. \end{cases}$$

Then we have that $\mathrm{Lip}(U_n) < c$ for any $n \in \mathbb{N}$. By the assumption, regarding $U_n \in U(C(\mathbb{T}) \otimes B)$, we have that $[U_n]_1 = \widetilde{\Delta}_\tau(U_n) = 0$ in $\tau(K_0(B))$.

Let B_n, $n \in \mathbb{N}$ be an increasing sequence of matrix subalgebras of B with $1_{B_n} = 1_B$ and $\overline{\bigcup B_n} = B$. Since $(U_n)_n \in U((C(\partial E) \otimes B)_\infty)$, slightly modyfying U_n, we obtain an increasing sequence $m_n \in \mathbb{N}$, $n \in \mathbb{N}$ and $U'_n \in U(C(\partial E) \otimes (B'_{m_n} \cap B))$ such that $m_n \nearrow \infty$, $(U'_n)_n = (U_n)_n$, $U'_n(0,0) = 1$, and $\mathrm{Lip}(U'_n) < c$. Since $B'_{m_n} \cap B$ has the unique tracial state $\tau|_{B'_{m_n} \cap B}$, it follows that $[U'_n]_{K_1(B'_{m_n} \cap B)} = \widetilde{\Delta}_{\tau_{B'_{m_n} \cap B}}(U'_n) = \widetilde{\Delta}_{\tau_B}(U'_n) = \widetilde{\Delta}_{\tau_B}(U_n) = 0$, then we obtain $\widetilde{U}_n \in U(C([0,1]^2) \otimes (B'_{m_n} \cap B))$, $n \in \mathbb{N}$ such that $\widetilde{U}_n|_{\partial E} = U'_n$ and $\mathrm{Lip}(\widetilde{U}_n) < c'$. Then we have that $(\widetilde{U}_n)_n \in (C([0,1]^2) \otimes B)_\infty$. Since $\widetilde{U}_n|_{\partial E} = U'_n$, $n \in \mathbb{N}$, slightly modyfying $\widetilde{U}_n$ on ∂E, we obtain $y_n \in U(C([0,1]^2) \otimes B)$, $n \in \mathbb{N}$ and $\varepsilon > 0$ such that $(y_n)_n = (\widetilde{U}_n)_n$, $y_n|_{\partial E} = U_n$, and $\mathrm{Lip}(y_n) < c' + \varepsilon$ for any $n \in \mathbb{N}$. ∎

As in the proof of Proposition 2.2 in [33], the embedding $I(k, k+1) \subset_{\text{unital}} (A_\infty)_{\alpha_\infty}$ obtained in Lemma 4.3 implies the following Lemma.

Lemma 5.5. *Let A be a a unital separable projectionless C^*-algebra which has a unique tracial state and absorbs the Jiang-Su algebra tensorially. Suppose that A has property (SI) and α is an automorphism of A with the weak Rohlin property. Then there exists a unital embedding of $\mathcal{Z}$ into $(A_\infty)_{\alpha_\infty}$.*

proof of Theorem 5.3. Let $\mathcal{Z}_k$, $B^{(i)}$, and $\Phi^{(i)}$, $i=0,1$, be the projectionless C^*-algebra, the UHF algebras, and the unital embeddings in the proof of Proposition 3.5.

Because of $\mathcal{Z}\otimes B^{(i)}\cong B^{(i)}$, the Rohlin property of $\sigma\otimes\mathrm{id}_{B^{(i)}}\in\mathrm{Aut}(\mathcal{Z}\otimes B^{(i)})$, and

$$\Delta_{\tau_{\mathcal{Z}\otimes B^{(i)}}}(u_n\otimes 1_{B^{(i)}})=0,\ \text{in } \mathbb{R}/\tau(K_0(B^{(i)})),\ n\in\mathbb{N},$$

applying Proposition 5.1, we obtain ${V_n}^{(i)}\in U(\mathcal{Z}\otimes B^{(i)})$, $i=0,1$, $n\in\mathbb{N}$ such that $({V_n}^{(i)})_n\in(\mathcal{Z}\otimes B^{(i)})_\infty$ and

$$\begin{aligned}&({V_n}^{(i)}\sigma\otimes\mathrm{id}_{B^{(i)}}({V_n}^{(i)})^*)_n=(u_n\otimes 1_{B^{(i)}})_n,\ i=0,1,\\ &\tau_{\mathcal{Z}\otimes B^{(i)}}\circ\log(V_n^{(i)}\sigma\otimes\mathrm{id}_{B^{(i)}}(V_n^{(i)})^*u_n^*\otimes 1_{B^{(i)}})=0,\ n\in\mathbb{N}.\end{aligned}$$

By the following argument, we would like to obtain a path of unitaries $\widetilde{v}_n$ in $\mathcal{Z}\otimes\mathcal{Z}_k$ with endpoints $\Phi^{(i)}(V_n^{(i)})\in U(\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})$, $i=0,1$ which satisfies $\widetilde{v}_n(t)\sigma\otimes\mathrm{id}_{B^{(0)}\otimes B^{(1)}}\ (\widetilde{v}_n(t)^*)\approx u_n\otimes 1_{B^{(0)}\otimes B^{(1)}}$ for any $t\in[0,1]$.

Set

$$\begin{aligned}U_{n,1}&=\Phi^{(0)}(V_n^{(0)})^*\Phi^{(1)}(V_n^{(1)}),\\ W_n&=U_{n,1}\sigma\otimes\mathrm{id}_{B^{(0)}\otimes B^{(1)}}(U_{n,1})^*,\ n\in\mathbb{N}.\end{aligned}$$

Then it follows that $(U_{n,1})_n\in(\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})_\infty$, $(W_n)_n=1_{(\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})_\infty}$, and, by (1) in Proposition 5.2, $\tau_{\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)}}\circ\log(W_n)$

$$\begin{aligned}&=\tau\circ\log(\Phi^{(1)}(V_n^{(1)}\sigma\otimes\mathrm{id}_{B^{(1)}}(V_n^{(1)})^*u_n^*\otimes 1_{B^{(1)}})\\ &\quad\cdot\Phi^{(0)}(V_n^{(0)}\sigma\otimes\mathrm{id}_{B^{(0)}}(V_n^{(0)})^*u_n^*\otimes 1_{B^{(0)}})^*)\\ &=\sum_{i=0,1}(-1)^{1-i}\tau\circ\log(\Phi^{(i)}(V_n^{(i)}\sigma\otimes\mathrm{id}_{B^{(i)}}(V_n^{(i)})^*)u_n^*\otimes 1_{B^{(i)}})=0,\end{aligned}$$

for any $n\in\mathbb{N}$.

Since $(U_{n,1})_n\in(\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})_\infty$, there exist $\widetilde{U}_n\in U(C([0,1])\otimes\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})$, $n\in\mathbb{N}$ be such that $\widetilde{U}_n(0)=1$, $\widetilde{U}_n(1)=U_{n,1}$, $(\widetilde{U}_n)_n\in(C([0,1])\otimes\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})_\infty$, and $\mathrm{Lip}(\widetilde{U}_n)<\pi+\varepsilon$ for some $\varepsilon>0$. Define $\widetilde{T}_n^{(j)}\in U(C([0,1])\otimes\mathcal{Z}\otimes B^{(0)}\otimes B^{(1)})$, j, $n\in\mathbb{N}$ by

$$\widetilde{T}_n^{(j)}=\widetilde{U}_n\,\mathrm{id}_{C([0,1])}\otimes\sigma^j\otimes\mathrm{id}_{B^{(0)}\otimes B^{(1)}}(\widetilde{U}_n)^*,$$

$\widetilde{T}_n^{(0)}=1$. Note that

$$\widetilde{T}_n^{(j)}\,\mathrm{id}\otimes\sigma\otimes\mathrm{id}(\widetilde{T}_n^{(j-1)})^*=\widetilde{T}_n^{(1)},\quad j,n\in\mathbb{N}.$$

By $(\widetilde{U}_n)_n \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$, $(W_n)_n = 1$, $\tau \circ \log(W_n) = 0$, and $\mathrm{Lip}(\widetilde{U}_n) < \pi + \varepsilon$, we have that $(\widetilde{T}_n^{(j)})_n \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$, $(\widetilde{T}_n^{(j)}(1))_n = 1$, $\tau\circ\log(\widetilde{T}_n^{(j)}(1)) = j\tau\circ\log(W_n) = 0$, and $\mathrm{Lip}(\widetilde{T}_n^{(j)}) < 2(\pi+\varepsilon)$, $j \in \mathbb{N}$. Then, by Lemma 5.4, we obtain a constant $c > 0$ and $y_n^{(j)} \in U(C([0,1]^2) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})$, $j \in \mathbb{N}$ such that $(y_n^{(j)})_n \in (C([0,1]^2) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$,

$$y_n^{(j)}(0,t) = 1, \quad y_n^{(j)}(1,t) = \widetilde{T}_n^{(j)}(t),\ t \in [0,1]$$

$$y_n^{(j)}(s,1) = \exp(\log(\widetilde{T}_n^{(j)}(1)s), \quad y_n^{(j)}(s,0) = 1,\ s \in [0,1],$$

and $\mathrm{Lip}(y_n^{(j)}) < c$, $n \in \mathbb{N}$.

By the Rohlin property of $\sigma \otimes \mathrm{id}_{B^{(0)}\otimes B^{(1)}}$ we obtain $p_m^{(l)} \in P(\mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})$ and $z_m^{(l)} \in U(\mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})$, l, $m \in \mathbb{N}$ such that $(p_m^{(l)})_m \in (\mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$, $(z_m^{(l)})_m = 1$, and

$$\sum_{j=0}^{k^l-1} (\mathrm{Ad}\, z_m^{(l)} \circ \sigma \otimes \mathrm{id})^j (p_m^{(l)}) = 1.$$

Set $y_n^{(j)}(s)(t) = y_n^{(j)}(s,t)$, $s, t \in [0,1]$, j, $n \in \mathbb{N}$,

$$\begin{aligned} \widetilde{\sigma}_m^{(l)} &= \mathrm{id}_{C([0,1])} \otimes (\mathrm{Ad}\, z_m^{(l)} \circ \sigma \otimes \mathrm{id}_{B^{(0)}\otimes B^{(1)}}), \quad l,\ m \in \mathbb{N}, \\ \widetilde{W}'_{l,m,n} &= \sum_{j=0}^{k^l-1} \widetilde{T}_n^{(j)} \cdot (\widetilde{\sigma}_m^{(l)})^{j-k^l} (y_n^{(k^l)}(j/k^l))^* \cdot (\widetilde{\sigma}_m^{(l)})^j (1_{C([0,1])} \otimes p_m^{(l)}). \end{aligned}$$

Since $((\widetilde{\sigma}_m^{(l)})^j\, (1_{C([0,1])} \otimes p_m^{(l)}))_m \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$, $j = 0,1,\ldots,k^l-1$ are mutually orthogonal projections, we have that $(\widetilde{W}'_{l,m,n})_m$ is a unitary and obtain $\widetilde{W}_{l,m,n} \in U(C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})$, l, m, $n \in \mathbb{N}$ such that $(\widetilde{W}_{l,m,n})_m = (\widetilde{W}'_{l,m,n})_m$. By the definition of $\widetilde{W}'_{l,m,n}$, we have that

$$\|(\widetilde{W}_{l,m,n})_m \cdot (\mathrm{id}_{C([0,1])} \otimes \sigma \otimes \mathrm{id}(\widetilde{W}_{l,m,n})^*)_m - (\widetilde{T}_n^{(1)})_m\| < c/k^l, \quad l,\ n \in \mathbb{N}.$$

Since $(\widetilde{T}_n^{(j)})_n$, $((\widetilde{\sigma}_m^{(l)})^{j-k^l} (y_n^{(k^l)}(j/k^l)))_n$, and $(\widetilde{\sigma}_m^{(l)})^j (1_{C([0,1])} \otimes p_m^{(l)})_m \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$, and $\|1 - W_n\| \to 0$, we obtain a slow increasing sequence l_n, $n \in \mathbb{N}$ and a fast increasing sequence $m_n \in \mathbb{N}$, $n \in \mathbb{N}$ such that $l_n \nearrow \infty$, $m_n \nearrow \infty$,

$$(\widetilde{W}_{l_n,m_n,n})_n \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty,$$

$$k^{2l_n}\|1 - W_n\| \to 0, \quad \|\widetilde{W}_{l_n,m_n,n}\, \mathrm{id} \otimes \sigma \otimes \mathrm{id}(\widetilde{W}_{l_n,m_n,n})^* - \widetilde{T}_n^{(1)}\| < c/k^{l_n}.$$

Set

$$\widetilde{V}'_n = \widetilde{W}^*_{l_n,m_n,n} \widetilde{U}_n \in U(C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)}), \quad n \in \mathbb{N}.$$

Then it follows that $(\widetilde{V}'_n)_n \in C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$ and

$$\begin{aligned} &(\widetilde{V}'_n\, \mathrm{id}_{C([0,1])} \otimes \sigma \otimes \mathrm{id}_{B^{(0)}\otimes B^{(1)}} (\widetilde{V}'_n)^*)_n \\ &= (\mathrm{Ad}\, \widetilde{W}^*_{l_n,m_n,n}(\widetilde{T}_n^{(1)} \cdot \mathrm{id} \otimes \sigma \otimes \mathrm{id}(\widetilde{W}_{l_n,m_n,n}) \cdot \widetilde{W}^*_{l_n,m_n,n}))_n = 1, \end{aligned}$$

Since $\|1-\widetilde{T}_n^{(j)}(1)\| \le j\|1-W_n\| \to 0$, and $\|1-y_n^{(k^{l_n})}(i/k^{l_n},1)\| \le \|1-\widetilde{T}_n^{(k^{l_n})}(1)\|$, it follows that

$$\begin{aligned}\widetilde{W}'_{l_n,m_n,n}(1) &= \sum_{j=0}^{k^{l_n}-1} \widetilde{T}_n^{(j)}(1)(\widetilde{\sigma}_{m_n}^{(l_n)})^{j-k^{l_n}}(y_n^{(k^{l_n})}(j/k^{l_n},1))^*(\widetilde{\sigma}_{m_n}^{(l_n)})^j(p_{m_n}^{(l_n)}) \\ &\approx_{\delta_n} 1.\end{aligned}$$

where $\delta_n = 2k^{2l_n}\|1-W_n\|$, $n \in \mathbb{N}$, and then we have that

$$(\widetilde{V}'_n(1))_n = (U_{n,1})_n = (\Phi^{(0)}(V_n^{(0)})^*\Phi^{(1)}(V_n^{(1)}))_n.$$

Define

$$\widetilde{V}_n(t) = \Phi^{(0)}(V_n^{(0)})\widetilde{V}'_n(t), \quad t \in [0,1].$$

Then we have that $(\widetilde{V}_n)_n \in (C([0,1]) \otimes \mathcal{Z} \otimes B^{(0)} \otimes B^{(1)})_\infty$

$$(\widetilde{V}_n(i))_n = (\Phi^{(i)}(V_n^{(i)}))_n,\ i = 0,1,$$

and that $(\widetilde{V}_n\, \mathrm{id}_{C([0,1])} \otimes \sigma \otimes \mathrm{id}_{B^{(0)}\otimes B^{(1)}}(\widetilde{V}_n)^*)_n$

$$\begin{aligned} &= (1_{C([0,1])} \otimes \Phi^{(0)}(V_n^{(0)}\sigma \otimes \mathrm{id}_{B^{(0)}}(V_n^{(0)})^*))_n \\ &= (1_{C([0,1])} \otimes u_n \otimes 1_{B^{(0)}\otimes B^{(1)}})_n.\end{aligned}$$

Slightly modyfying $\widetilde{V}_n$ at the end points, we obtain $\widetilde{v}_n \in U(\mathcal{Z} \otimes \mathcal{Z}_k)$ such that $(\widetilde{v}_n)_n = (\widetilde{V}_n)_n$,

$$\widetilde{v}_n(i) = \Phi^{(i)}(V_n^{(i)}),\ i = 0,1, \quad (\widetilde{v}_n\sigma \otimes \mathrm{id}_{\mathcal{Z}_k}(\widetilde{v}_n)^*)_n = (u_n \otimes 1_{\mathcal{Z}_k})_n.$$

Finally, we obtain $(v_n)_n \in \mathcal{Z}_\infty$ which corresponds to $\widetilde{v}_n \in \mathcal{Z}\otimes\mathcal{Z}_k$ and satisfies $(v_n\sigma(v_n^*))_n = (u_n)_n$, by the following. By Lemma 5.5 and $\mathcal{Z}_k \subset_{\mathrm{unital}} \mathcal{Z}$, we obtain a unital embedding $\Psi : \mathcal{Z} \otimes \mathcal{Z}_k \hookrightarrow \mathcal{Z}^\infty$ such that $\sigma_\infty \circ \Psi = \Psi \circ \sigma \otimes \mathrm{id}_{\mathcal{Z}_k}$ and $\Psi(a \otimes 1_{\mathcal{Z}_k}) = a \in \mathcal{Z} \subset \mathcal{Z}^\infty$, $a \in \mathcal{Z}$. Let $F_n \subset \mathcal{Z}^1$, $n \in \mathbb{N}$ be an increasing sequence of finite subsets of $\mathcal{Z}^1$ and $\varepsilon_n > 0$, $n \in \mathbb{N}$ a decreasing sequence such that $\overline{\bigcup F_n} = \mathcal{Z}^1$, $\varepsilon_n \searrow 0$,

$$\|[\widetilde{v}_n, x \otimes 1_{\mathcal{Z}_k}]\| < \varepsilon_n, \quad x \in F_n.$$

$$\|\widetilde{v}_n\sigma \otimes \mathrm{id}_{\mathcal{Z}_k}(\widetilde{v}_n)^* - u_n \otimes 1_{\mathcal{Z}_k}\| < \varepsilon_n,$$

It follows that $\|[\Psi(\widetilde{v}_n), x]\| = \|\Psi([\widetilde{v}_n, x \otimes 1_{\mathcal{Z}_k}])\| < \varepsilon_n$, $x \in F_n$, and

$$\begin{aligned}\Psi(\widetilde{v}_n)\sigma_\infty(\Psi(\widetilde{v}_n)^*) &= \Psi(\widetilde{v}_n\sigma \otimes \mathrm{id}_{\mathcal{Z}_k}(\widetilde{v}_n)^*) \\ &\approx_{\varepsilon_n} \Psi(u_n \otimes 1_{\mathcal{Z}_k}) = u_n.\end{aligned}$$

Denote by $v_{n,p} \in U(\mathcal{Z})$, $p \in \mathbb{N}$ components of $\Psi(\widetilde{v}_n) \in U(\mathcal{Z}^\infty)$, then we obtain an increasing sequence $p_n \in \mathbb{N}$, $n \in \mathbb{N}$ such that

$$(v_{n,p_n}\sigma(v_{n,p_n})^*)_n = (u_n)_n, \quad (v_{n,p_n})_n \in \mathcal{Z}_\infty.$$

Define $v_n = v_{n,p_n}$. This completes the proof. ∎

proof of Theorem 1.3. By using the stability of the following form instead of Proposition 4.3 [21] and by the Evans-Kishimoto intertwining argument in the proof of Theorem 5.1 [21] we can give the proof, where we omit the detail.

Corollary 5.6. *Suppose that $\sigma \in \mathrm{Aut}(\mathcal{Z})$ has the weak Rohlin property. For any finite subset F of $\mathcal{Z}^1$ and $\varepsilon > 0$, there exists a finite subset G of $\mathcal{Z}^1$ and $\delta > 0$ satisfying that: for any $u \in U(\mathcal{Z})$ with $\|[u, y]\| < \delta$, $y \in G$, there exist $v \in U(\mathcal{Z})$ and $\lambda \in \mathbb{T}$ such that*

$$\|v\sigma(v)^* - \lambda u\| < \varepsilon, \quad \|[v, x]\| < \varepsilon, \ x \in F.$$

Proof. For $u_n \in U(\mathcal{Z})$, $n \in \mathbb{N}$ with $(u_n)_n \in \mathcal{Z}_\infty$, set $\lambda_n = \exp(-2\pi\sqrt{-1}\,\Delta_{\tau_{\mathcal{Z}}}(u_n)) \in \mathbb{T}$. Since $\Delta_{\tau_{\mathcal{Z}}}(\lambda_n u_n) = 0 \in \mathbb{R}/\tau(K_0(\mathcal{Z}))$, by the above theorem we obtain $v_n \in U(\mathcal{Z})$, $n \in \mathbb{N}$ such that $(v_n)_n \in \mathcal{Z}_\infty$ and

$$(v_n\sigma(v_n)^*)_n = (\lambda_n u_n)_n.$$

Assume that there exist a finite subset F of $\mathcal{Z}^1$ and $\varepsilon > 0$ satisfying that: For any finite subset G of $\mathcal{Z}^1$ and $\delta > 0$ there exists $u \in U(\mathcal{Z})$ with $\|[u, y]\| < \delta$, $y \in G$ such that if $v \in U(\mathcal{Z})$ and $\lambda \in \mathbb{T}$ satisfy $\|v\sigma(v)^* - \lambda u\| < \varepsilon$ then $\|[v, x]\| \geq \varepsilon$ for some $x \in F$. This contradicts the above statement. ∎

Acknowledgments. The author would like to thank his advisor Akitaka Kishimoto for many valuable discussions, and he also would like to thank Hiroki Matui for the preprint [23] and useful advices.

References

[1] D. Archey, *Crossed product C^*-algebras by finite group actions with the projection free tracial Rokhlin property*, arXiv: 0902.3324.

[2] A. Connes, *Outer conjugacy classes of automorphisms of factors*, Ann. Scient. Ec. Norm. Sup. (4) **8** (1975), 383-419.

[3] M. Dadarlat and W. Winter, *Trivialization of $C(X)$-algebras with strongly self-absorbing fibres*, Bull. Soc. Math. France 136 (2008), no. 4, 575-606.

[4] M. Dadarlat, N. C. Phillips, and A. S. Toms, *A direct proof of $\mathcal{Z}$-stability for AH algebras of bounded topological dimension*, arXiv: 0806.2855.

[5] K. R. Davidson, *C^*-algebras by Example.* Fields Inst. Monographs, Vol. 6, Amer. Math. Soc. Providence, 1996.

[6] M. Dadarlat, and W. Winter, *On the K-theory of strongly self-absorbing C^*-algebras*, Math. Scand. 104 (2009), no. 1, 95-107.

[7] G. A. Elliott, *An invariant for simple C^*-algebras,* Canadian Mathematical Society. 1945-1995 3 (1996), 61-90.

[8] G. A. Elliott, and A. S. Toms, *Regularity properties in the classification program for separable amenable C^*-algebras*, Bull. Amer. Math. Soc. 45 (2008), no.2, 229-245.

[9] D. E. Evans and A. Kishimoto, *Trace scaling automorphisms of certain stable AF algebras*, Hokkaido Math. J. 26 (1997), 211-224.

[10] G. Gong, X. Jiang, and H. Su, *Obstructions to $\mathcal{Z}$-stability for unital simple C^*-algebras*, Canad. Math. Bull. 43 (2000), no. 4, 418-426.

[11] P. de la Harpe and G. Skandalis, *Déterminant associé à une trace sur une algébre de Banach*, Ann. Inst. Fourier (Grenoble) 34 (1984), 241-260.

[12] I. Hirshberg and W. Winter, *Rokhlin actions and self-absorbing C^*-algebras.* Pacific J. Math. 233 (2007), no. 1, 125-143.

[13] M. Izumi, *The Rohlin property for automorphisms of C^*-algebras*, Mathematical Physics in Mathematics and Physics (Siena, 2000), 191-206, Fields Inst. Commun., 30, Amer. Math. Soc., Providence, RI, 2001.

[14] M. Izumi, *Finite group actions on C^*-algebras with the Rohlin property. I*, Duke Math. J. 122 (2004), 233-280.

[15] M. Izumi, *Finite group actions on C^*-algebras with the Rohlin property. II*, Adv. Math. 184 (2004), 119-160.

[16] M. Izumi and H. Matui, *$\mathbb{Z}^2$-actions on Kirchberg algebras*, preprint, arXiv:0902.0194.

[17] H. Lin, *Inductive Limits of Subhomogeneous C^*-algebras with Hausdorff Spectrum*, preprint, arXiv:0809.5273.

[18] X. Jiang and H. Su, *On a simple unital projectionless C^*-algebra,* Amer. J. Math. 121 (1999), no. 2, 359-413.

[19] T. Katsura and H. Matui, *Classification of uniformly outer actions of $\mathbb{Z}^2$ on UHF algebras*, Adv. Math. 218 (2008), 940-968. arXiv:0708.4073.

[20] A. Kishimoto, *The Rohlin property for automorphisms of UHF algebras*, J. Reine Angew. Math. 465 (1995), 183-196.

[21] A. Kishimoto, *Unbounded derivations in AT algebras*, J. Funct. Anal. 160 (1998), 270-311.

[22] H. Matui, *Classification of outer actions of $\mathbb{Z}^N$ on $\mathcal{O}_2$*, Adv. Math. 217 (2008), 2872- 2896. arXiv:0708.4074.

[23] H. Matui, *$\mathbb{Z}$-actions on AH algebras and $\mathbb{Z}^2$-actions on AF algebras*, arXiv: 0907.2474.

[24] H. Nakamura, *Aperiodic automorphisms of nuclear purely infinite simple C^*-algebras*, Ergodic Theory Dynam. Systems 20 (2000), 1749-1765.

[25] H. Osaka and N. C. Phillips, *Stable and real rank for crossed products by automorphisms with the tracial Rokhlin property*, Ergodic Theory Dynam. Systems 26 (2006), no. 5, 1579-1621.

[26] G. K. Pedersen, *C^*-algebras and their automorphism groups*, London Mathematical Society Monographs, 14. Academic Press, Inc. 1979.

[27] M. V. Pimsner, *Ranges of traces on K_0 of reduced crossed products by free groups*, Operator algebras and their connections with topology and ergodic theory (Busteni, 1983), 374-408, Lecture Notes in Math., 1132, Springer, Berlin, 1985.

[29] M. Rørdam, *The stable and the real rank of $\mathcal{Z}$-absorbing C^*-algebras,* International J. Math. 15 (2004), no. 10, 1065-1084.

[30] M. Rørdam and W. Winter, *The Jiang-Su algebra revisited,* to appear in J. Reine Angew. Math. arXiv: 0801.2259.

[31] Y. Sato, *Certain aperiodic automorphisms of unital simpl projectionless C^*-algebras,* to appear in Internat. J. Math. arXiv: 0807.4761.

[32] Y. Sato, *A generalization of the Jiang-Su construction*, preprint, arXiv: 0903.5286.

[33] A. S. Toms and W. Winter. *$\mathcal{Z}$-stable ASH algebras,* Canad. J. Math. 60 (2008), no. 3, 703-720. arXiv:math/0508218

[34] W. Winter, *Localizing the Elliott conjecture at strongly self-absorbing C^*-algebras - with an appendix by Huaxin Lin -*, arXiv: 0708.0283.

[35] W. Winter. *Strongly Self-absorbing C^*-algebras are $\mathcal{Z}$-stable*, arXiv: 0905.0583.